

constant amount of active ingredient in each dose, to give predictable relief from occasional or non-persistent constinution.

BEFORE TAKING THIS MEDICINE: Do not take if:

· your abdomen is tender to the touch or hurts when you move vou have sharp or persistent stomach pain

Consult your doctor or pharmacist if:

there is no bowel movement after three days

- laxatives are needed every day, or abdominal pain persists
- · you accidentally take too many tablets
- - if you have been told that you have an intolerance to some sugars as this product contains sucrose.

HOW TO TAKE SENOKOT MAX STRENGTH TABLETS:

Senokot Max Strength Tablets are to be taken by mouth only. Adult and children over 12: take 1 tablet at night. Do not take more than 1 tablet in

You may experience temporary mild stomach pains. If you have these

any 24hr period. Children 12 and under: consult your doctor.

Senokot Max Strength Tablets usually act within 8-12 hours. The tablets should be taken with a glass of water. Dose can be repeated on a daily basis until bowel action is restored. SIDE EFFECTS

or any other symptoms after taking this product, tell your doctor or pharmacist.

(S) FURTHER INFORMATION

Each tablet contains standardised senna equivalent to 15mg total sennosides in a base containing calcium phosphate, maize starch,

lactose and magnesium stearate. Keep out of reach and sight of children. Don't store above 30°C. Store in the original package. Do not use after the expiry date shown

on the pack (EXP month/year) UK PL 00063/0118 Senokot and (†) are trade marks Last revised April 2011

