

# What Original Andrews Salts Do

Original Andrews Salts quickly and effectively relieve:

-  Indigestion
-  Excess Acid
-  Symptoms of Over-Indulgence
-  Upset Stomach
-  Constipation

The market authorisation holder is GlaxoSmithKline Consumer Healthcare, Brentford, TW8 9GS, U.K. and all enquiries should be sent to this address.

The manufacturer is GlaxoSmithKline SA, 28806 Alcala de Henares, Madrid, Spain.

Information was last revised in August 2008.

Andrews is a registered trade mark of the GlaxoSmithKline group of companies.

## How to use Original Andrews Salts

See inside label for how to open and dosage instructions.

Check before you take this medicine if you need a laxative dose every day or you have persistent abdominal pain (tummy ache).

## Further information

Effervescent powder containing the active ingredients Sodium Hydrogen Carbonate 22.6% w/w, Citric Acid 19.5% w/w and Magnesium Sulphate 17.4% w/w. Also contains 2.1 g of sucrose per 5 ml spoonful.



Keep out of the reach and sight of children.

Use within 3 months of opening. Check inner seal is intact before first use. Contents sold by weight and may settle after packaging. Store below 25°C and away from strong odours.


BN:

9 42  
0 6 20 12

Do not use  
after end:



250 g



5 000347 068003 >

PL 00071/5000R L393004/05  
11000000003341

Peel here but do  
not remove



## How to take Original Andrews Salts

### How to open:

Lift end of lid. Check inner seal is intact before first use. Break inner seal carefully.

### How to take:

**Adults:** *For upset stomach, indigestion and over-indulgence*, measure one level spoonful (5 ml spoonful) into a glass of water and drink. Take as needed, up to a maximum of 4 times a day.

*For constipation*, measure two level spoonfuls (two 5 ml spoonfuls) into a glass of water and drink. Take before breakfast or at bedtime.

**Children over 3 years:** Use half the adult dose.

**Not suitable for children under 3 years.**

Close the lid after use.

**If you take too much you may get diarrhoea. If this happens, reduce the dose and the effect should go away.**

## Check before you take Original Andrews Salts:

- **Ask your doctor or pharmacist:**
- **if you are taking and prescribed medicines; particularly tetracycline antibiotics (e.g. minocycline)**
- **if you are pregnant**
- **if you are on a controlled sodium diet. Each 5 ml spoonful contains 319 mg of sodium**
- **if you have high blood pressure, heart or kidney disease and avoid frequent use.**

## Tell your doctor or pharmacist:

- **if you need a laxative dose every day or you have persistent abdominal pain (tummy ache)**
- **if symptoms persist**
- **if you experience any unwanted effects.**