

PLEASE READ THIS LEAFLET CAREFULLY BEFORE YOU START TO USE THIS MEDICINE. IT CONTAINS IMPORTANT INFORMATION, WHICH YOU SHOULD KEEP FOR REFERENCE.

Manufacturer of Deep Relief

The licence holder and manufacturer of **Deep Relief** is The Mentholatum Company Limited, 1 Redwood Avenue, Peel Park Campus, East Kilbride, G74 5PE, Scotland.

Mode of Action

Deep Relief is a clear colourless gel containing Ibuprofen (5%) and Levomenthol (3%). The dual action formulation allows rapid and concentrated absorption of the active ingredients through the skin, direct to the point of pain.

Ibuprofen is one of a group of medicines called non-steroidal anti-inflammatory drugs which provide effective sustained pain relief and reduce both inflammation and swelling.

Levomenthol acts as a counter-irritant providing an immediate soothing sensation to calm the pain.

The result of this combination is immediate pain relief which lasts for hours.

Each tube contains 15, 30 or 50g; also available in a pump action dispenser containing 50g.

Ingredients

Active ingredients:

Ibuprofen

50mg/g

Levomenthol

30mg/g

Other ingredients:

propylene glycol, carbomer, di-isopropanolamine, ethanol and water.

Indications

Relief of rheumatic pain, muscular aches, pains and swellings such as strains, sprains and sports injuries.

Precautions in use

1. Do not use the product if:

- · You are allergic to ibuprofen, aspirin or any of the ingredients in Deep Relief.
- You have asthma and know that aspirin or non-steroidal anti-inflammatory drugs may bring on an asthma attack, rhinitis, itching or rashes.
- · You are pregnant or think you may be pregnant.

2. Do not apply to the following areas:

- Lips, nostrils, eyes, genital or anal areas, mucous membranes, broken or inflamed skin.
- If this occurs accidentally, wash away with plenty of clean water.

3. Consult a doctor or pharmacist before use in the following

- History of allergic symptoms caused by medicines or cosmetics (eg rashes/itching etc.).
- If you have a kidney problem.
- · If you are taking aspirin or any other medicine.
- You are a mother currently breast feeding.



4. Precautions prior to use

- Do not exceed the stated dosage.
- Do not use on children under 12 years old.
- · If a serious condition develops or any unwanted effects are seen consult your doctor.
- This medicine is for external use only. Do not take orally.

5. Directions for use

- For adults and children over 12 years, apply the gel as a thin layer over the affected area. For each application, use about 1-4cm if using the 30 or 50g size and use 4-10cm if using the 15g size.
- Gently massage until the gel is absorbed. You will notice a tingling sensation.
- Repeat these steps up to 3 times a day but no more. Do not repeat more frequently than every 4 hours.
- Unless treating the hands always wash your hands after use. Replace the cap tightly.
- Do not exceed the stated dose.

6. Cautions after use

- If you swallow the gel, contact your doctor or nearest casualty department immediately.
- · If the pain or swelling continues after using the gel for two weeks consult your doctor.
- Do not cover the skin with bandages, plasters or any other dressing. It does not matter if you cover the skin with your clothes.

7. Side effects

- Sometimes unwanted effects can occur in some people.
- A slight redness and tingling may occur where you have applied the gel. This is not a cause
- However, if this gets worse or if you experience any of the following, stop using the medicine immediately and tell your doctor: unexplained wheezing, shortness of breath, skin rash, itching, bruising, or facial swelling.
- Stomach pains or indigestion may rarely be noticed.
- . If this is a concern, or you experience any other effects, you should speak to your doctor or pharmacist.

8. Storage and handling

- Keep all medicines out of the reach of children.
- Close the cap tightly and store in a cool place (not above 25°C).
- Do not use the product after the expiry date (printed on the container).



