

**PACKAGE LEAFLET:  
INFORMATION FOR THE USER**

**Dioralyte Relief Blackcurrant**

**Dioralyte Relief Raspberry**

**6.0g/580mg/350mg/300mg,**

**Powder for oral solution**

Pre-cooked rice powder, sodium citrate, sodium chloride, potassium chloride

**SANOFI** 

**Is this leaflet hard to see or read?  
Phone 0845 372 7101 for help**



**Read all of this leaflet carefully because it contains important information for you.**

This medicine is available without prescription. However, you still need to use Dioralyte Relief Blackcurrant/Raspberry carefully to get the best results from it.

- **Keep this leaflet.** You may need to read it again.
- **Ask your pharmacist** if you need more information or advice.
- **You must contact a doctor** if your symptoms worsen or do not improve after using this medicine for 36 hours.
- **If you notice any side effects** not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What Dioralyte Relief is and what it is used for
2. Before you take Dioralyte Relief
3. How to take Dioralyte Relief
4. Possible side effects
5. How to store Dioralyte Relief
6. Further information

**1. What Dioralyte Relief is and what it is used for**



The name of your medicine is Dioralyte Relief Blackcurrant/Raspberry (called Dioralyte Relief throughout this leaflet). Dioralyte Relief works by replacing the water and salts lost from your body when you have diarrhoea. It helps the watery stools return to normal.

It contains pre-cooked rice powder and salts (these are sodium citrate, sodium chloride and potassium chloride). The powder is mixed with water before taking.

**2. Before you take Dioralyte Relief**



**Do not take Dioralyte Relief and talk to your doctor or pharmacist if:**

- × You are allergic (hypersensitive) to pre-cooked rice powder, sodium citrate, sodium chloride, potassium chloride or any of the other ingredients of Dioralyte Relief (see Section 6: Further information)
- Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue
- × Your bowel is partially or fully blocked (called 'intestinal obstruction')
- × You have severe kidney or liver disease

- × You have something called 'phenylketonuria' (a rare illness passed on in families)
  - × The child taking the medicine is under 3 months old
- Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Dioralyte Relief.



**Take special care with Dioralyte Relief**  
**Check with your doctor or pharmacist before taking this medicine if:**

- ▲ You have something called an 'electrolyte imbalance' (unusual levels of salts in your body)
  - ▲ You are on a low potassium or low sodium diet
  - ▲ **The child taking the medicine is under 1 year old**
- If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Dioralyte Relief.



**Taking other medicines**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines you buy without a prescription, including herbal medicines. This is because Dioralyte Relief can affect the way some other medicines work. Also some medicines can affect the way Dioralyte Relief works.

**Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine if you are pregnant or breast-feeding. This includes if you are planning to get pregnant or think that you might be pregnant.

**Important information about some of the ingredients of Dioralyte Relief**

Dioralyte Relief contains a source of phenylalanine (aspartame). This may be harmful to people with something called 'phenylketonuria'.

Dioralyte Relief contains 2.5 vol % ethanol (alcohol). One dose contains 191 mg of ethanol, which is about the same amount of alcohol as 4 ml of beer or 1.7 ml of wine. This can be harmful to people with a drink problem. Also take this into account if you are pregnant, breast-feeding or giving this to a child, have liver disease or epilepsy.

**3. How to take Dioralyte Relief**


If you are not sure how to take this medicine, check with your doctor or pharmacist

**Preparing this medicine**

Dioralyte Relief must be mixed with water before taking.

- Prepare just before the medicine is going to be taken. Do not make in advance. If you are giving it to a baby over the course of the day, you can keep the drink in a refrigerator for up to 24 hours
- **Only** dissolve the powder in water. Do not dissolve it in anything else
- Use fresh drinking water for adults and children over 1 year old. If fresh water is not available, use water which has been freshly boiled and cooled
- Always use water which has been freshly boiled and cooled for babies aged 3 months to 1 year





- 1 Empty the contents of 1 sachet into a glass
  - 2 Add 200 ml (7 fl oz) of water
  - 3 Stir until the powder dissolves
- When your medicine is fully dissolved, the drink will look milky

Do not use more or less than 200 ml of water. If you do, the drink will become too weak or too strong. It will not work as well. You may also upset the balance of salts in your body.

#### How many sachets to take

##### Adults (including the elderly) and children over 1 year

- One sachet after each loose stool
- You can take up to 5 sachets each day, for up to 3 to 4 days. Do not take more than 5 sachets in 24 hours

##### Babies aged 3 months to 1 year (to be given under the guidance of a doctor)

- The amount depends on how much the baby weighs. The usual dose is 150-200 ml of the drink for each kg of the baby's weight, each day. This can be given over 24 hours
- Give half the drink during the first 8 hours and the other half during the next 16 hours
- Ask your doctor or pharmacist if you are not sure how much to give
- If your baby is being sick, you can give 5-10 ml (one or two teaspoons) every 5 minutes. This can be gradually increased until they are able to drink normally.

**If you take more Dioralyte Relief than you should** Dioralyte Relief is not normally harmful if you take more than you should. However, if large amounts are taken, you should talk to a doctor or go to a hospital straight away. This is particularly important if it is a child or baby. Take the medicine pack with you.

##### If you forget to take your Dioralyte Relief

Do not take a double dose to make up for a forgotten dose. Take the next dose when you need to.



If you have any further questions on the use of this product, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, Dioralyte Relief can cause side effects, although not everybody gets them. However, if you notice anything unusual, talk to your doctor or pharmacist. You should not normally have any side effects.

**Stop taking Dioralyte Relief and see a doctor or go to a hospital straight away if:**

- You have an allergic reaction. The signs may include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)  
By reporting side effects you can help provide more information on the safety of this medicine.

#### 5. How to store Dioralyte Relief

Keep out of the sight and reach of children.

Do not use Dioralyte Relief after the expiry date which is stated on the carton and sachet after EXP. The expiry date refers to the last day of that month.

Store in a dry place below 25°C.

The drink should be made up just before it is going to be taken. Throw away any remaining drink after 1 hour (or 24 hours if stored in a refrigerator).

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### 6. Further information

##### What Dioralyte Relief contains

- The active substances are pre-cooked rice powder, sodium citrate, sodium chloride and potassium chloride
- The other ingredients are hypromellose, aspartame, blackcurrant flavour 501017 or raspberry flavouring 611417E, purified water and ethanol 96%

##### What Dioralyte Relief looks like and contents of the pack

Dioralyte Relief is available in pack sizes of 6. Packs of 6 may contain:

- 6 sachets of one flavour
- 4 sachets of blackcurrant flavour and 2 of raspberry (Blackcurrant & Raspberry)
- 4 sachets of raspberry flavour and 2 of blackcurrant (Raspberry & Blackcurrant)

Each sachet contains enough powder to make a drink up with 200 ml (7 fl oz) of water.

#### Marketing Authorisation Holder and Manufacturer

##### Marketing Authorisation Holder

Sanofi  
One Onslow Street  
Guildford Surrey GU1 4YS, UK  
Tel: 0845 372 7101  
email: [uk-medicalinformation@sanofi.com](mailto:uk-medicalinformation@sanofi.com)

##### Manufacturer

Sanofi-Aventis Sp. z o.o.,  
52, Lubelska Str., 35-233 Rzeszów, Poland

This leaflet does not contain all the information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

**This leaflet was last revised in 01/2014.**

© Sanofi, 1995-2014