

Cherry Flavour

Ultra Chloraseptic® Anaesthetic Throat Spray

Please read this leaflet carefully before you start using your medicine, as it contains important information for you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If a side effect occurs and it seems serious to you, or if you notice a side effect not listed in this leaflet, tell your doctor or pharmacist.

1. What Ultra Chloraseptic® Anaesthetic Throat Spray is and what it is used for

Ultra Chloraseptic® Anaesthetic Throat Spray is used to relieve sore throat pain.

The product contains the active ingredient benzocaine which belongs to the group of medicines called local anaesthetics. It has a fast action and works by numbing the pain of a sore throat. The product comes in the form of a spray. It has a swivel head applicator to help deliver the local anaesthetic direct to the throat.

2. Before you use Ultra Chloraseptic® Anaesthetic Throat Spray

DO NOT use this spray if

- you have difficulty in breathing, noisy breathing or severe difficulty in swallowing. Talk to your pharmacist or doctor if this applies to you
- you know you are allergic to benzocaine or any of the other ingredients (see Section 6, Further Information)
- you have been told that you have a rare blood condition called methaemoglobinaemia
- you are pregnant or breast-feeding

DO NOT use this medicine for children under 6 years of age.

This medicine contains a small quantity of alcohol (less than 0.006 units per spray) to dissolve the other ingredients. Contains propylene glycol - this may cause skin irritation.

If your sore throat is severe or prolonged, or accompanied by fever, headache, or feeling sick, consult a doctor.

In this leaflet:

1. What Ultra Chloraseptic® Anaesthetic Throat Spray is and what it is used for
2. Before you use Ultra Chloraseptic® Anaesthetic Throat Spray
3. How to use Ultra Chloraseptic® Anaesthetic Throat Spray
4. Possible side effects
5. How to store your medicine
6. Further information


3. How to use Ultra Chloraseptic® Anaesthetic Throat Spray

Ultra Chloraseptic® Anaesthetic Throat Spray is for use by adults and teenagers, and in children aged from 6 years when supervised by an adult.

The product is designed so that the medication can be sprayed directly to the back of the throat.

The following instructions will help you to get the best results from your spray:


- Remove the cap from the bottle, and **rotate the spray head around** by half a circle so that the nozzle extends beyond the bottle (see diagram below).



- If this is a new bottle, or you have not used it since your last sore throat, **activate the pump by spraying 3 times away from your face into a sink**. This will also help you to get a feel for the pump action.
- Hold the bottle in your hand with your index (first) finger on the pump, ready to spray.

3. How to use (CONTINUED)

- When you are ready, open your mouth a little and **point the spray nozzle towards the back of your throat**. The nozzle should be just inside your mouth (see diagram below). This will help the spray get to its target at the back of the throat.
- **Hold your breath and then press the pump to spray.** Do not exceed the dosage on the labelling.



- It may be easier the first few times to stand in front of the mirror whilst spraying the product.

Spray Dosage:

DO NOT spray into the eyes. If you accidentally spray into your eyes, rinse them thoroughly under water.

For Adults and Teenagers:

- Hold breath and spray 3 times to the back of your throat.
- Repeat every 2-3 hours if required.
- Do not use more than 8 times a day.

For Children 6-12 years:

- Use only under adult supervision.
- Do not use in a child who is unable to hold their breath whilst spraying.
- Hold breath and spray once to the back of the throat.
- Repeat every 2-3 hours if required.
- Do not use more than 8 times a day.

DO NOT use for more than 3 days in a row. If symptoms do not improve, consult your pharmacist or doctor.

You should experience temporary numbness in your throat after using the spray. This indicates that the product is working. **Avoid eating or drinking as long as the numbness lasts.**

If you spray more than you should, and you feel drowsy, you must contact a doctor immediately.

4. Possible side effects

Allergic reactions have been reported occasionally with benzocaine. Although very unlikely, if your breathing is affected after using the spray, or if your face or mouth become swollen, stop using the spray and seek immediate medical advice.

Methaemoglobinaemia (a very rare blood condition) has been reported with benzocaine use.

If you experience any side effect whilst taking this medicine, tell your doctor or pharmacist.

5. How to store your medicine

Keep out of the reach and sight of children.

Do not use the spray after the Expiry date which is shown on the top of the carton and on the base of the spray bottle.

6. Further information

What Cherry Flavour Ultra Chloraseptic® Anaesthetic Throat Spray contains

The spray contains Benzocaine 0.71% w/v (1mg per spray) as the active ingredient. It also contains: ethanol, macrogol 300, propylene glycol, glycerol, cetylpyridinium chloride, saccharin sodium, sodium dihydrogen phosphate dihydrate, sodium hydroxide, lactic acid, flavouring (cherry, cream, and levomenthol) and purified water.

What Cherry Flavour Ultra Chloraseptic® Anaesthetic Throat Spray looks like and contents of pack

The spray is a clear colourless to straw coloured solution, presented in a glass bottle with a spray nozzle under a plastic cap. Each bottle contains 15ml. The nozzle is used to target the spray to the back of the throat.

The marketing authorisation for Cherry Flavour Ultra Chloraseptic® Anaesthetic Throat Spray is held by: Prestige Brands (UK) Limited, 3 Scotlands Drive, Farnham Common, Slough, Berkshire SL2 3ES, UK. The product is manufactured at: Wägener & Co GmbH, Poststrasse 24, 49525 Lengerich, Germany. 785958001 Version 3.3

This leaflet is dated October 2010.

© Prestige Brands (UK) Limited 2010

® Chloraseptic is a registered trademark.

To get this leaflet in large print, Braille or as an audio CD, call the RNIB Medicine Leaflet line on **0800 198 5000** and quote **PL 18259-0002**.